

Strengthening Teaching Competences in Higher Education in Natural and Mathematical Sciences

WORKSHOP NOVI SAD MARCH 21.-22., 2022



KATARÍNA TRNKOVÁ

MATEJ BEL UNIVERSITY, FACULTY OF NATURAL SCIENCES, BANSKÁ BYSTRICA, SLOVAKIA



#### EDUCATION AND PROFESSIONAL CAREER

M.Sc. degree in study programme General Ecology

**Ph.D.** (in external form) of study programme **Ecology**, on the Faculty of E Sciences, Technical University Zvolen

2001–2013 work position as laboratory diagnostics specialist (public health monitoring of water, indoor and outdoor air quality) in The Regional Authority of Public Health (RAPH) in BB, Department of Medical Microbiology: (in 2006 the **specialization** in study programme

**Investigation methods in Microbiology and Biology of the Environment,** Slovak Medical University in Bratislava

2014 – till now: Assistant Professor, Faculty of Science, Matthias Belius University, Banská Bystrica Department of Environment, but closely cooperated with Department of Biology and Ecology



I was participate in the teaching in the **previous study programs** Environmental safety and Environmental management, Ecology and ecosystem protection till 2020/2021

#### of subjects such as

Health and Environment Environmental Epidemiology The Global Environmental Problems Bioindications in the Environment Biological and Chemical Terrorism Safety Introduction to the study of Environmental Sciences

Monitoring of Environmental Compartment Environmental Loads and Brownfields Remediation of Environmental Loads



#### Faculty of Natural Sciences, MBU in B. Bystrica obtained accreditation in field

#### **Ecological and Environmental Sciences**

new Study program (2021/2022): Environmental Biology (bachelor, master, doctoral degree)

Characteristics of the study program:

"From molecules to ecosystems" - to know the basic organizational levels of life (genes, cells, organisms and ecosystems and their relationships with the environment),

to learn about the processes that shaped nature in the past and affect it today,

find out what and how life-threatening systems on Earth threaten

you will understand how to solve the environmental problems of water, soil, air, even with the effective help of living organisms - bioindication


## **ENVIRONMENTAL BIOLOGY SUBJECTS THAT I LEAD**

- Environmental Health
- Epidemiology
- Molecular Ecology
- Microbiology
- Global Problems of Environment
- Abiotic components of the Environment part II. (Water and Air)

practice exercises, excursions and internships presentation skills (Student scientific conference, Researchers night, Open day of Faculty NS, etc.)

Our students will use the acquired knowledge and skills in research, nature protection, the environment and other professions Cooperation with, government organizations, institutions, and the third sector











1

\*\*

#### WE TEACH DIFFERENTLY, WE TEACH FOR PRACTICE



#### SEMESTRAL GROUP WORK CONNECTING THEORY WITH PRACTICE



learning, practical training and cooperation with businesses (contracting authority Nestlé, Veolia - local waterworks company), institutions (SHMÚ Slovak hydrometeorological institute), and the third sector (Ekopolis foundation)

The idea of our team was to proposed **artificial pond** supplied with **rainwater drained** from the **roof** of an industrial building in Sewage treatment plants area using climatic, pedological, biotic technical-material and economy calculation data

National competition of student teams - theme "For water" (we won 2. place)



Stredoslovenská vodárenská prevádzková spoločnosť, a.s.





## **PROJECT IDEA AND GOALS**

BIODIVERSITY INCREASING – to creat conditions for increasing the natural biodiversity of the environment




## WORKFLOW

- 1. introduction of project goals, working meeting with contracting
- 2. work schedule, division of partial project tasks into groups of students
- 3. soil sampling, laboratory analyzes, collection and interpretation of data, project promotion
- 4. second working meeting summarization of documents and data to create a technical proposal project of the pond
- 5. risk processing of project, preparing information panels design and content
- 6. project finalization, and presentation











### DESIGN OF TECHNICAL PROJECT OF THE POND




FIG. 3 Scheme of implementation of the pond, legend: 1. roof as a source of rainwater, 2. mechanical pre-treatment, 3. wetland - filter part of the pond, 4. pond, 5. overflow, 6. wet meadow

## **PROJECT COSTS CALCULATION**

Work / material	Price
Technical work (excavation of the pit, modification of the pond)	€ 008
Material (pond foil, geotextile, river gravel, sewer pipe, roof drainage catcher)	1884.20 €
Bacteria Starter	20 €
Plants	625€
Information panels	400 €
Sum	3729.20 €

## **HOW INCREASE BIODIVERSITY?**





Veľký a malý vodný cyklus

### PREVÓDU addčia sekopolisi Prejavy nestabiného počasia, striedanie obdobia sucha s obdobim prudkých prívalových daždov, či významný nárast rozlohy

Prejavy nestabilného počasia, striedanie obdobia sucha s obdobím prudkých privalových dažďov, či významný nárast rozlohy nepriepustných plôch z dôvodu výstavby parkovisk, nákupných centier, komunikácií, domov je veľkým problémom súčasnosti. Pri lokálnych bůrkach dochádza ku zaťaženiu mestskej kanalizácie privalovou vodou zo spevnených plôch, čo potom spôsobuje vzostup vody v potokoch a riekach, dochádza k záplavám.



Εκοιγοιέπουο ταιδέπέ printupy κ αστράσί sù παέπου atternatiou atueso dopinion w παδύλη τοιποίοητη πότομοη, του χύης αραστείατι, Elosopatineou zaudole pristup a princie obiae optimeta, note vigualiza isolocetta a consolatione attudi va notimy element cenavaj adaptable prisobatenta sa reprizavjun esposion zimery kilin. Teo optimeta optimetaliju proste eleñena, ao noti prividante zelenjon pôch na priva valenej inflažitutativa (zelené optimeta) – domona a vijinicante zelenjon pôch a priva, and a era inflicientare na citata a renytipi relacional, zelendo zelendo zelendo zelendo zelendo zelendo za prividante zelendo zelendo.

stiech a zelených lasád budov, výsadba stromov a nahrádzanie spevnených plách trávnívní, budovanie daždových záhrad; využité stojaných aj tečúcich vôd v urbanizovanom prostredí, obnova mokradi

2) prvky modrej infraštruktúry – revitalizácia vodných tokov, zvyšovanie zadržiavania vody ej aj vomej krajine, zvyšovanie priepustnosti terénu v mestách





A C SPÔSOBY VYUŽ VDO zavlažovanie, kúre

a umyvanie au

strecha : makrad 4. jazierko 5. prepad

VLHKÁ LÚKA pre vyššiu biodiverzitu by mala obsahova maximum autochtónnych (pôvodných) hygrofytnýci



### WE HAVE LEARNED A LOT, WE HAVE DONE A LOT 😇




http://www.humanisti.sk/view.php?nazevclanku=studenti-univerzity-mateja-bela-v-banskej-bystrici-budu-h%beadat-riesenie-pre-cistiare%f2-odpadovych-v%f4d-v-rakytovciach&cisloclanku=2017110031

#### WHAT NEXT?

I will introduce you with the topic of paleoecology, I will explain the basic concepts and its application,

I will tell you how I connect research activities with the teaching and assignment of final theses and the involvement of students in research.



# PALYNOLOGY - USE TODAY TO KNOW THE PAST

## **TERMINOLOGY / GLOSSARY**

- Palynology is the "study of dust" (from <u>Greek palunō</u>, "strew, sprinkle"and <u>-logy</u>) or "particles that are strewn"
- A classic palynologist analyses particulate samples collected from the air, from water, or from deposits including sediments of any age
- The condition and identification of those particles, organic and anorganic, give the palynologist clues to the life, environment, and energetic conditions that produced them
- Palynology is an interdisciplinary science and is a branch of earth science (geology or geological science) and biological science (biology), particularly botany
- Stratigraphical palynology is a branch of micropalaeontology and paleobotany, which studies fossil palynomorphs from the Precambrian to the Holocene (to the Anthropocene 😌)

#### **APPLICATIONS**

- Biostratigraphy and geochronology. Geologists use palynological studies in biostratigraphy to correlate strata and determine the relative age of a given bed, horizon, formation or stratigraphical sequence
- Palaeoecology and climate change palynology can be used to reconstruct past vegetation (land plants) and marine and freshwater phytoplankton communities, and so infer past environmental (palaeoenvironmental) and palaeoclimatic conditions
- Palynology also allows scientists to infer the climatic conditions from the vegetation present in an area thousands or millions of years ago. This is a fundamental part of research into climate change
- Organic palynofacies studies, which examine the preservation of the particulate organic matter and palynomorphs provides information on the depositional environment of sediments and depositional palaeoenvironments of sedimentary rocks
- Limnology studies. Freshwater palynomorphs and animal and plant fragments, including the (green algae, pollen, chironomids head capsules, diatoms, cladocera, ostracods) can be used to study past lake levels and long term climate change
- Allergy studies. Studies of the geographic distribution and seasonal production of pollen, can help sufferers of allergies such as hay fever
- Archaeological palynology examines human uses of plants in the past. This can help determine seasonality of site occupation, presence or absence of agricultural practices or products, and 'plant-related activity areas' within an archaeological context
- Melissopalynology: the study of pollen and spores found in honey

### WHAT ARE WE GOING TO TALK ABOUT?


 Palynology (Aerobiology) use at present – "Pollen information service in Slovakia"

analysis of recent pollen grain

 Pollen analysis – reconstruction of past environmental conditions – "Tracking Environmental Change Using Lake Sediments"

analysis of fosil nollon grain






## **TERMINOLOGY**

- Aerobiology (from <u>Greek ἀήρ</u>, aēr, "air"; <u>βίος</u>, bios, "<u>life</u>"; and <u>-λογία</u>, <u>-logia</u>) is a branch of biology that studies organic particles, such as bacteria, fungal spores, very small insects, pollen grains and viruses, which are passively transported by the air
- Aerobiologists have traditionally been involved in the measurement and reporting of airborne pollen and fungal spores as a service to allergy sufferers (Hayfever affects 1 in 5 people with symptoms like: itching mouth and throat, swelling of mouth, tongue and lips, hoarseness, numbness, swelling of eyelids, digestion problems)

## **POLLEN INFORMATION SERVICE IN SLOVAKIA**

- The negative trend of an increase of allergic diseases, especially pollen allergy, has been an incentive for physicians, botanists and aerobiologists to initiate building of a network of PIS monitoring stations in Slovakia working since 2006 on public health authorities responsibility
- Based on laboratory analyzes of the pollen of the allergenic relevant gere reports are processed
- Pollen reports are publish for both, gand a forecast for the next week national electronic and print media)



### POLLINATION SPECIES OR ALLERGENIC RELEVANT SPECIES

A six basic groups of plants producing pollen allergens, to which the majority of polinotic are sensitive, have been identified in Europe:

- birch (Betula) + related species alder, hazel, hornbeam (Alnus, Corylus, Carpinus) beech (Fagus), chestnut(Castanea) a oak (Quercus)
- grasses (Poaceae) + wheat (Gramineae)
- Oleaceae + ash (Fraxinus), olive (Olea) + maple (Acer), walnut (Juglandaceae) and (Platanaceae)
- weed plants (Artemisia) + (Ambrosia), (Asteraceae), Chenopodiaceae, Plantaginaceae)
- (*Parietaria officinalis*) + nettle (*Urtica dioica*)
- (*Cupressaceae*) + related species (*Taxaceae*) + *Cryptomeria japonica*

## POLLEN PERIODS IN CENTRAL EUROPE

In Central Europe to split the season into:

• spring season - dominated by a pollen of trees, especially birch (Betula) and hazel (Corylus)

• summer season - the dominant allergens are grasses

• and autumn - with the dominance of high-bred weeds, especially Artemisia, and in recent years also Ambrosia, even though their considerable production of pollen is still in the late summer

## POLLEN PERIODS IN MEDITERRANEAN AREA

In the Mediterranean area there are three pollen seasons:

- A low winter pollen season (December March) with pollens from trees such as Cypress and Juniper, Hazel, Mimosa and Birch
- A high spring-summer pollen season (April July), dominated by the pollination of Grasses, some members of the nettle family and olive trees. Between March and May, *Platanus* pollen (such as the American sycamore) may effect some people in Mediterranean areas
- A summer- autumn season (August October) involves a second flowering of the nettle family and pollination of herbaceous plants, such as mugwort Artemisia and plants such as beet, sugar beet, and chard, and wild spinach Chenopodiaceae)

### **POLLEN MONITORING METHODS** - METHODOLOGY OF AEROBIOLOGY


- Volumetric method standard monitoring method of PIS allows to express the concentration of biological particles in volume units of air (an important value for expressing the possible patient load) Measurements of the pollen content in the air are the fundament of every pollen forecast
- Pollen traps (pollen sampler) are used in aerobiology for this purpose and answer the question of what was and is in the air
- continuous and volumetric (Hirst type, e.g. Burkard, Lanzoni) flow rate of 10 L/min sampling height between 15-20 m above ground


location distant from high buildings or other obstacles, trees and other local pollen sources

a description of the surrounding (urban/rural area, valley/hill side, vegetation) should be available



## POLLEN TRAP BURKARD






Adhesive tapes as capturing surfaces in Burkard sampling


Adhesives: vaseline and silicon oil are recommended depending on the climatic conditions

### SAMPLES PREPARING






Tapes is cutting - each sample is divided in days and bi-hourly sections **Imbedding medium:** most often gelvatol or glycerine jelly are used **Staining:** mostly basic fuchsin or saffranine



## **POLLEN COUNTING**

- Pollen is determined and counted on the slide by the light microscope compound light microscope magnification 400 x resolution (numerical aperture of the objective) not below 0.65
- Counting method: In the qualitative and quantitative analysis of the daily sample, the method of horizontal strips of touching fields is used at least 10% of the sample has to be analyzed (this means 4 longitudinal bands, 12 transverse (vertical) bands (a conversion factor has to be applied depending on the microscope and the method used to receive daily averages ad daily concentration (pollen per m<sup>3</sup> air) corresponding to the average daily content of pollen grains and spores
- Those values are used for the pollen forecast, the graphs, models and many more services

#### And so I learned to know the pollen spectrum ... 7 years practice with pollen monitoring 😌







## POLLEN REPORTS AND FORECAST

#### https://www.polleninfo.org/country-choose.html



## **POLLEN INFORMATION SERVICE IN EU**

• Pollen Information Service - monitoring of biological allergens in the air

In Europe, peer monitoring is provided through interconnected national networks of PIS stations

European Aeroallergen Network initiative in building of

A central European pollen database with European Aeroallergen network Server (EANS) started activity in 1988, Vienna

So, the isolated pollen services of each member of European countries have been linked to a pan-European network

At present, around 300 pollen stations from around this area are contributing to the European pollen database

https://www.polleninfo.org/country-choose.html

### PALYNOLOGY AND ITS USE FOR RECONSTRUCTION OF PAST ENVIRONMENTAL CONDITIONS

Paleolimnological research presented through our national projects focused to climatic evolution of the Central Europe in post glacial period.

Why can we use pollen in research?

Bioindicators

Pollen grains preserved in sediments are very resistant to environmental influences, which allows them to remain preserved and intact for a long period of time. The ability of long inertia is enabled by their outer shell called **exina** (for pollen grains) and **exospores** (for spores) (Moore et al., 1991). The chemical compound that allows the exine of pollen grains to withstand mechanical, physical and chemical influences is called **polenin**. In spores, this compound is referred to as sporenine

#### **OUR PALEOLIMNOLOGY RESEARCH**

latest ongoing project named

 TAMARA - acronym (Reading in natural archives: ThousAnds of years of environmental history and cliMAte change Recorded in Alpine lakes of the Ukrainian Carpathians)

The project aims to provide a detailed reconstruction of climate and environmental changes during the Holocene in this mountain region

Specifically, the research will be focusing on the following topics:

1. Reconstruction of the mean July air temperature trends over the Holocene

2. Identification of the timing and range of long-term human-induced effects on the lakes and alpine landscape

3. Defining the timing and intensity of stressors affecting the lakes during last century (acidification, eutrophication)

#### **OUR PALEOLIMNOLOGY RESEARCH**

 "A 250 year history of human impacts on a landscape of the Banská Štiavnica mining territory: a palaeolimnological reconstruction using biotic and abiotic indicators from lake sediments"

The aim: (mining, multi-proxy approach, catchment vegetation changes, reservoir ontogeny, heavy metals, Anthropocene)

 "Deglaciation and postglacial climatic development of the High Tatras recorded in the lake sediments"

The aim: climatic evolution of the Central Europe in post glacial period

Pollen analysis is very valuable paleoclimate proxy and it was used for paleoenvironmental reconstructions of the High Tatras landscape, in this project, pollen analysis were used as a proxy to identify changes in plant communities at the transition from clay (glacial) and to organic (post-glacial) sedimentation

http://www.geo.sav.sk/en/depovyt-apvv-15-0292/





### MINNING RESHAPED THE RELIEF

Hundreds of years of the ore mining reshaped the relief andha s left a legacy of abandoned surface mines,

subsidence pits, mine adits, abandoned heap of metallurgical s lag, dumps, and reservoirs and rests of the canal system







These structures became an integral part of the landscape and persist until the present day

Existence of the unique, man-modified geomorphology has been one of the reasons to establish the Protected Landscape Area Štiavnické vrchy in 1979

In 1993, Banská Štiavnica together with the remains of the hydro-energetic system were inscribed in the World Cultural and Natural Heritage List of UNESCO

## **RESERVOIRS AND RESTS OF THE CANAL SYSTEM**

- Because the fundamental and extensive changes in the Banská Štiavnica region have occurred back in the past before scientific observation was possible, palaeolimnological methods have to be employed to study the timing, severity and extent of human impacts
- At the present time, 30 reservoirs of the former hydro-energetic system are preserved and serve mostly for recreation and angling. They potentially contain deposits that represent an important and unique archive of reservoir and watershed dynamics
- To get an overview of the natural and anthropogenic events recorded in the sediments we have to look back into the past by disentangling the changes in sub-fossil organisms and sediment characteristics (i.e. 'proxy' data)

#### SCIENTIFIC GOALS

1) To reconstruct changes in vegetation cover in the reservoir watersheds as a response to intensity of human impacts: from heavy mining practices to natural and managed reforestation

2) To determine the anthropogenic impacts on past sediment delivery from the deforested watersheds during the mining period and to identify changes in sediment transfer related to with the development of vegetation

3) To figure out (based on the chemical composition of the sediments) the impact of mine wastes, especially heavy metals, on the environment over time, and in connection with the findings to assess the potentially toxic risk for humans e.g. via food chain (drinking water, fish)

4) To find a stratigraphic marker from which the beginning of Anthropocene in the northern part of the Austro-Hungarian Monarchy could be identified

5) To reconstruct changes of lake ecosystems over time: how aquatic organisms responded to extreme conditions prevailing in the period of the biggest boom of mining (water-level fluctuation, extensive influx of allochthonous material, toxic effects of pollutants) and how the succession advanced after mining activities ended?

We will track the development trajectories of two model reservoirs under different management conditions over the past ~ 100 years


## **SEDIMENT EXTRACTION**

- Year 2015 and 2016 steel coreworker working on a hydraulic principle
- Sonar survey selecting the site with the largest sediment thickness
- Length of sedimentation cores (V. Richňava 184 cm, Vindšachta 120 cm, Rozgrund 420 cm)





#### LABORATORY ANALYSIS







Determination skills, evaluation and interpretation of results, to train presentation skills on Student research conference competitions - winner of scholarship







